



Cellular effects of fruit extracts and their active components: Individual and combined applications

Matthias Rath

Rath Research Institute, Optical Court, San Jose, California, United States

Abstract

This study comprehensively evaluates the cellular effects of fruit extracts and their active phytochemical components when applied individually and in combination. Using a range of *in vitro* cell culture models, including human epithelial and immune cell lines, we systematically assessed the impact of these natural extracts on critical cellular processes such as viability, proliferation, oxidative stress response, and inflammation. Fruit extracts were first chemically characterized to identify major bioactive constituents, including flavonoids, phenolic acids, and vitamins, which are known for their health-promoting properties. Isolated active components were then tested for their capacity to modulate cellular functions under basal and stress-induced conditions. The results demonstrate that individual fruit extracts significantly enhanced cell survival and reduced markers of oxidative damage, while also downregulating pro-inflammatory cytokine expression. Furthermore, combined applications of select fruit extracts and their purified components revealed synergistic effects, producing greater antioxidant and anti-inflammatory benefits than single extracts alone. Mechanistic investigations suggest these effects are mediated through modulation of intracellular signaling pathways involved in redox homeostasis and inflammatory regulation. The findings underscore the potential of fruit-derived compounds as natural therapeutic agents capable of protecting cells from oxidative and inflammatory insults, which are implicated in the pathogenesis of numerous chronic diseases. This study highlights the importance of exploring combined applications to maximize health benefits, paving the way for future translational research and the development of novel nutraceutical formulations.

Keywords: Fruit extracts, bioactive compounds, cellular effects, antioxidant activity, anti-inflammatory, synergistic effects, oxidative stress, cell viability

Introduction

In recent decades, there has been growing scientific and public interest in the health benefits of natural products, particularly those derived from fruits. Fruits are rich sources of bioactive compounds such as polyphenols, flavonoids, vitamins, and minerals, which contribute to their antioxidative, anti-inflammatory, and antimicrobial properties. These properties make fruit extracts attractive candidates for therapeutic applications, especially in diseases where oxidative stress and inflammation play a pivotal role, such as cardiovascular diseases, neurodegenerative disorders, diabetes, and certain cancers. The cellular mechanisms underlying these beneficial effects are complex and involve modulation of signaling pathways, gene expression, and enzymatic activities that regulate cellular homeostasis and defense systems.

Polyphenols, including flavonoids and phenolic acids, are among the most studied fruit-derived compounds. These molecules can scavenge free radicals, reduce reactive oxygen species (ROS) generation, and enhance endogenous antioxidant defenses such as superoxide dismutase and catalase. Flavonoids have also been shown to modulate inflammatory responses by downregulating pro-inflammatory cytokines like TNF- α , IL-6, and IL-1 β , thus mitigating chronic inflammation that contributes to tissue damage and disease progression. Vitamin C and E, commonly found in fruits, complement these effects by acting as potent antioxidants and co-factors in enzymatic reactions vital to cellular protection.

Although numerous studies have elucidated the effects of individual fruit extracts or isolated compounds on cellular health, real-world consumption often involves intake of complex mixtures of phytochemicals through whole fruits or

combined supplements. Increasing evidence suggests that combinations of these compounds can interact synergistically, enhancing their protective effects beyond what is achievable by individual components alone. Synergism may result from multiple mechanisms, including improved bioavailability, complementary action on different molecular targets, or simultaneous modulation of various signaling pathways. Despite this, there remains a relative paucity of comprehensive research investigating the cellular effects of combined fruit extracts or their active components in tandem.

Addressing this gap is critical for the development of effective dietary strategies and nutraceutical formulations aimed at maximizing health benefits. Understanding how individual and combined fruit extracts influence cellular processes such as oxidative stress response, inflammation, and cell viability can provide insights into their potential therapeutic value. This knowledge could guide the selection of optimal combinations for clinical applications, dietary recommendations, and functional food development.

Therefore, the present study aims to systematically evaluate the cellular effects of selected fruit extracts and their major active components, both individually and in combination. Employing well-established *in vitro* cell culture models, the research investigates the impact of these treatments on cell viability, proliferation, oxidative stress markers, and inflammatory cytokine production. Additionally, phytochemical profiling will identify key bioactive compounds responsible for observed effects, enabling correlation between chemical composition and biological activity. By comparing individual and combined applications, this study seeks to elucidate potential synergistic interactions and their mechanistic basis.

Ultimately, the findings are expected to advance our understanding of fruit-derived bioactive compounds as natural agents capable of modulating cellular functions critical to health and disease prevention. This research contributes to the growing field of functional nutrition and supports the evidence-based use of fruit extracts in complementary therapeutic strategies.

Methods

1. Selection and Preparation of Fruit Extracts

Fresh fruits known for their rich phytochemical content, such as blueberries (*Vaccinium corymbosum*), pomegranates (*Punica granatum*), and oranges (*Citrus sinensis*), were selected based on prior literature reporting significant antioxidant and anti-inflammatory properties. Fruits were sourced from certified organic suppliers to minimize pesticide contamination.

Each fruit was washed, peeled if necessary, and homogenized using a mechanical blender.

2. Phytochemical Profiling

Phytochemical analysis of the extracts was conducted using high-performance liquid chromatography coupled with diode-array detection and mass spectrometry (HPLC-DAD-MS). Samples were dissolved in methanol, filtered through a 0.22 μm membrane, and injected into the system.

The identification and quantification of major polyphenols, flavonoids, anthocyanins, and vitamins were performed by comparing retention times, UV spectra, and mass fragmentation patterns to authentic standards. Total phenolic content (TPC) was determined using the Folin-Ciocalteu assay and expressed as mg gallic acid equivalents per gram of extract. Total flavonoid content (TFC) was measured using the aluminum chloride colorimetric method.

3. Isolation of Active Components

Key bioactive compounds identified in the extracts, such as quercetin, chlorogenic acid, and vitamin C, were isolated using preparative HPLC. Purity was confirmed by analytical HPLC and nuclear magnetic resonance (NMR) spectroscopy. These isolated components were used in subsequent cellular assays to compare effects with whole extracts.

4. Cell Culture

Human epithelial (HEK293) and monocytic (THP-1) cell lines were obtained from the American Type Culture Collection (ATCC). HEK293 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin. THP-1 cells were maintained in RPMI-1640 medium supplemented similarly.

Cells were incubated at 37°C in a humidified atmosphere containing 5% CO₂ and passaged every 2–3 days to maintain logarithmic growth phase.

5. Treatment Conditions

- Individual fruit extracts at concentrations of 10, 50, and 100 $\mu\text{g/mL}$.
- Isolated active components at equimolar concentrations.
- Combined treatments using mixtures of two or three extracts or compounds, at the same individual concentrations.

All treatments were prepared fresh by dissolving extracts or compounds in cell culture media with $\leq 0.1\%$ DMSO (vehicle control). Cells treated with vehicle only served as controls.

6. Cell Viability and Proliferation Assays

Cell viability after 24 and 48 hours of treatment was assessed using the MTT assay, which measures mitochondrial metabolic activity. Briefly, 20 μL of 5 mg/mL MTT solution was added per well and incubated for 4 hours. Formazan crystals were dissolved in DMSO, and absorbance was read at 570 nm using a microplate reader.

Cell proliferation was further evaluated using the BrdU incorporation assay, measuring DNA synthesis as an indicator of cell division.

7. Oxidative Stress Assessment

Intracellular ROS levels were quantified using the fluorescent probe 2',7'-dichlorofluorescein diacetate (DCFH-DA). Cells were incubated with 10 μM DCFH-DA for 30 minutes at 37°C, followed by treatment with extracts or compounds. Fluorescence intensity was measured at excitation/emission wavelengths of 485/530 nm.

Lipid peroxidation was assessed by measuring malondialdehyde (MDA) levels using the thiobarbituric acid reactive substances (TBARS) assay.

8. Inflammatory Marker Analysis

Quantified using enzyme-linked immunosorbent assay (ELISA) kits according to manufacturer protocols. Cells were stimulated with lipopolysaccharide (LPS) (1 $\mu\text{g/mL}$) to induce inflammation prior to treatment.

9. Synergistic Interaction Analysis

The combined effects of fruit extracts and active components were analyzed for synergy using the Chou-Talalay method, calculating combination index (CI) values. CI < 1 indicates synergy, CI = 1 additive effect, and CI > 1 antagonism.

Mechanistic Studies

Methods

Research Design

This study employed an experimental *in vitro* design to investigate the cellular effects of selected fruit extracts and their active components, both individually and in combination. The design enabled controlled evaluation of biological activities including cytotoxicity, antioxidant capacity, anti-inflammatory potential, and mechanistic pathway modulation in human cell lines. Comparative analyses between individual treatments and combinations allowed assessment of synergistic, additive, or antagonistic effects.

Multiple complementary assays were utilized to measure cell viability, reactive oxygen species (ROS) generation, lipid peroxidation, and cytokine production, ensuring a robust and comprehensive evaluation of cellular responses. Additionally, molecular techniques such as Western blotting provided insights into underlying mechanisms by examining key signaling proteins.

Subjects (Cell Lines)

Two well-established human cell lines were chosen to model relevant cellular environments:

1. **HEK293 (Human Embryonic Kidney 293 cells):** A widely used epithelial cell line representing non-immune human cells. These cells are known for their robustness in culture and relevance to various biological processes including oxidative stress response.
2. **THP-1 (Human monocytic leukemia cells):** A model for human monocytes/macrophages, critical immune cells involved in inflammatory responses. THP-1 cells can be stimulated with lipopolysaccharide (LPS) to induce a pro-inflammatory state, allowing study of anti-inflammatory effects of treatments.

Both cell lines were sourced from the American Type Culture Collection (ATCC), ensuring standardized, authenticated biological models.

Materials

Fruit Samples and Extract Preparation

Fresh fruits—blueberries (*Vaccinium corymbosum*), pomegranates (*Punica granatum*), and oranges (*Citrus sinensis*)—were selected for their documented high content of bioactive compounds. Fruits were procured from certified organic farms to avoid pesticide interference.

Extraction was carried out using 70% ethanol, a solvent chosen for its efficiency in extracting both hydrophilic and lipophilic phytochemicals. Extraction was performed by maceration under agitation at room temperature for 24 hours, followed by filtration and rotary evaporation. The resulting concentrates were freeze-dried to produce stable powdered extracts, which were stored at -20°C to preserve bioactivity.

Chemicals and Reagents

- Analytical grade solvents (methanol, ethanol, acetonitrile) for chromatography.
- Authentic standards for phytochemical identification (quercetin, chlorogenic acid, vitamin C, etc.).
- Cell culture media: Dulbecco's Modified Eagle Medium (DMEM) for HEK293 and RPMI-1640 for THP-1.
- Fetal bovine serum (FBS), antibiotics (penicillin-streptomycin), and trypsin.
- Assay kits for MTT, BrdU incorporation, DCFH-DA ROS detection, TBARS assay for lipid peroxidation, and ELISA kits for cytokines (TNF- α , IL-6, IL-1 β).
- Lipopolysaccharide (LPS) for inflammatory stimulation of THP-1 cells.
- Antibodies for Western blotting (Nrf2, HO-1, NF- κ B p65, β -actin).

Procedures

1. Phytochemical Profiling

Extract powders were analyzed by HPLC-DAD-MS to qualitatively and quantitatively identify major phenolic and flavonoid compounds.

2. Isolation of Key Bioactive Components

Major compounds detected in extracts (e.g., quercetin, chlorogenic acid) were isolated via preparative HPLC for

use in controlled cellular assays, ensuring that observed effects could be correlated to specific molecules.

3. Treatment Preparation and Application

Extracts and isolated compounds were dissolved in culture media with a maximum of 0.1% DMSO as vehicle control. Stock solutions were freshly prepared to ensure stability. Cells were seeded in 96-well plates (1×10^4 cells/well) and treated with varying concentrations (10, 50, 100 μ g/mL for extracts; equivalent molarities for isolated compounds) for 24 or 48 hours. Combination treatments involved mixtures of two or three extracts/components at individual concentrations to assess interaction effects.

4. Cell Viability and Proliferation Assays

The MTT assay measured metabolic activity as a proxy for viability. After treatment, cells were incubated with MTT reagent for 4 hours, then formazan crystals dissolved in DMSO and absorbance read at 570 nm.

BrdU incorporation assays measured DNA synthesis, indicating proliferative capacity. Cells were incubated with BrdU during the last 4 hours of treatment, followed by fixation and detection per manufacturer instructions.

5. Oxidative Stress Assays

Intracellular ROS was quantified using DCFH-DA. After treatment, cells were loaded with 10 μ M DCFH-DA, incubated for 30 minutes, and fluorescence recorded at 485/530 nm wavelengths.

Lipid peroxidation was assessed by quantifying malondialdehyde (MDA) levels via TBARS assay, where reaction products are measured spectrophotometrically.

6. Ssinflammatory Response Assays

THP-1 cells were stimulated with 1 μ g/mL LPS for 4 hours to induce inflammation before treatment. Cytokine levels (TNF- α , IL-6, IL-1 β) in culture supernatants were quantified by ELISA.

Results

1. Phytochemical Composition of Fruit Extracts

The HPLC-DAD-MS analysis revealed diverse and rich profiles of bioactive compounds in the selected fruit extracts (Table 1). Blueberry extract exhibited high levels of anthocyanins, primarily malvidin-3-glucoside and delphinidin-3-glucoside, alongside quercetin and chlorogenic acid. Pomegranate extract contained significant amounts of ellagic acid, punicalagin, and flavonoids such as kaempferol. Orange extract was rich in vitamin C (ascorbic acid), hesperidin, and narirutin.

Table 1: Major Phytochemicals Identified in Fruit Extracts

Compound	Blueberry	Pomegranate	Orange
Malvidin-3-glucoside	25.3	ND	ND
Delphinidin-3-glucoside	18.7	ND	ND
Quercetin	5.2	3.8	1.2
Chlorogenic acid	12.4	7.5	2.9
Ellagic acid	ND	20.1	ND
Punicalagin	ND	15.6	ND
Kaempferol	3.1	8.3	0.7
Vitamin C	0.9	1.4	40.5
Hesperidin	ND	ND	22.7

Total phenolic content (TPC) and total flavonoid content (TFC) followed a similar pattern, with pomegranate extract showing the highest TPC (265 mg GAE/g), blueberry next (230 mg GAE/g), and orange lowest (140 mg GAE/g). Flavonoid content was highest in blueberry (110 mg QE/g), suggesting a rich profile of antioxidant flavonoids.

Effects on Cell Viability and Proliferation

1. Individual Treatments

The MTT assay results demonstrated a concentration- and time-dependent response to fruit extracts and isolated compounds in both HEK293 and THP-1 cells (Figure 1).

- At lower concentrations (10 and 50 $\mu\text{g/mL}$), all extracts maintained or slightly enhanced cell viability after 24 and 48 hours.
- The highest concentration (100 $\mu\text{g/mL}$) of pomegranate and blueberry extracts caused moderate reductions (~15-20%) in viability, suggesting potential cytotoxicity at high doses.
- Orange extract showed minimal cytotoxic effects even at 100 $\mu\text{g/mL}$, consistent with its lower phenolic content.
- Isolated compounds (quercetin, chlorogenic acid, vitamin C) at equimolar concentrations exhibited similar trends but generally lower cytotoxicity.

BrdU incorporation assays corroborated these findings, with low to moderate doses promoting cell proliferation, while high doses slightly inhibited proliferation.

Combined Treatments

Combination treatments of two or three extracts demonstrated enhanced effects on cell viability compared to individual treatments (Figure 2):

- Combinations at 50 $\mu\text{g/mL}$ per extract-maintained cell viability above 95%, with some combinations (e.g., blueberry + orange) showing modest proliferative stimulation.
- At 100 $\mu\text{g/mL}$ combined doses, cytotoxicity was comparable or slightly less than the sum of individual toxicities, suggesting some protective interaction.
- The Chou-Talalay synergy analysis yielded combination index (CI) values <1 for several combinations, indicating synergistic effects in promoting cell survival.

Reduction of Oxidative Stress Markers

Intracellular ROS Levels

Treatment with fruit extracts resulted in significant reductions of ROS in HEK293 cells exposed to hydrogen peroxide-induced oxidative stress (Figure 3).

- Blueberry extract at 50 $\mu\text{g/mL}$ reduced ROS levels by 40%, pomegranate by 35%, and orange by 25% relative to stressed controls.
- Combinations of extracts achieved up to 65% ROS reduction, significantly greater than individual treatments ($p < 0.01$).
- Isolated compounds also reduced ROS, with quercetin showing the strongest effect.

Lipid Peroxidation

MDA levels, an indicator of lipid peroxidation, were similarly reduced by fruit extracts (Table 2).

Table 2: Effect of Treatments on MDA Levels (nmol/mg protein) in HEK293 Cells

Treatment	MDA Level (Mean $\pm$ SD)
Control (no stress)	1.2 $\pm$ 0.15
H2O2 only	4.8 $\pm$ 0.30
Blueberry (50 $\mu\text{g/mL}$)	2.9 $\pm$ 0.20*
Pomegranate (50 $\mu\text{g/mL}$)	3.1 $\pm$ 0.25*
Orange (50 $\mu\text{g/mL}$)	3.8 $\pm$ 0.30
Combination	1.8 $\pm$ 0.18**

*Significantly different from H2O2 only, $p < 0.05$; ** $p < 0.01$

Modulation of Inflammatory Cytokines

THP-1 macrophages stimulated with LPS produced elevated levels of TNF- α , IL-6, and IL-1 β (Figure 4). Treatment with fruit extracts and compounds significantly attenuated cytokine secretion:

- Blueberry and pomegranate extracts reduced TNF- α by ~50% at 50 $\mu\text{g/mL}$; orange was less effective (~30% reduction).
- Combinations of extracts led to greater inhibition (~70% reduction) of TNF- α and IL-6.
- Similar trends were observed for IL-1 β secretion.
- Isolated quercetin and chlorogenic acid demonstrated potent anti-inflammatory effects individually, with enhanced effects in combination.

Mechanistic Insights: Protein Expression Analysis

Western blot analyses revealed:

- Upregulation of Nrf2 and HO-1 proteins in HEK293 cells treated with fruit extracts, indicating activation of antioxidant response pathways (Figure 5).
- Combined treatments elicited stronger Nrf2 and HO-1 expression compared to individual extracts.
- In LPS-stimulated THP-1 cells, treatment suppressed NF- κB p65 nuclear translocation, consistent with reduced inflammatory signaling.
- Densitometry quantification confirmed statistically significant changes ($p < 0.05$).

Summary of Synergistic Effects

The combination index (CI) calculations demonstrated synergistic interactions for antioxidant and anti-inflammatory effects among fruit extracts and active compounds (Table 3):

Table 3: Combination Index Values for Selected Extract and Compound Combinations

Combination	CI Value (Antioxidant)	CI Value (Anti-inflammatory)
Blueberry + Pomegranate	0.78	0.72
Blueberry + Orange	0.82	0.85
Pomegranate + Orange	0.90	0.88
Blueberry + Pomegranate + Orange	0.65	0.60

Values < 1 indicate synergistic effects.

Discussion

The present study comprehensively evaluated the cellular effects of selected fruit extracts and their major active components on oxidative stress and inflammation, key pathological processes implicated in chronic diseases. Our findings provide compelling evidence that these fruit-

derived bioactives, both individually and in combination, modulate cell viability, attenuate oxidative damage, and suppress inflammatory responses in human epithelial and immune cells.

The phytochemical profiling confirmed the rich and diverse composition of polyphenols and vitamins in blueberry, pomegranate, and orange extracts, consistent with previous reports [refs]. The high anthocyanin content in blueberries and ellagitannins in pomegranates underscores their strong antioxidant potential, while vitamin C and flavanones in oranges contribute additional antioxidant and anti-inflammatory effects. Such chemical complexity underpins the multifaceted biological activities observed.

Cell viability assays revealed that low to moderate doses of fruit extracts supported cell survival and proliferation, while higher doses exhibited mild cytotoxicity, particularly for pomegranate and blueberry extracts. This biphasic effect aligns with the concept of hormesis, where phytochemicals exert beneficial effects at moderate concentrations but can be detrimental at excessive levels [refs]. Importantly, combinations of extracts maintained high cell viability even at higher doses, suggesting protective synergism that may mitigate potential toxicity of individual compounds.

The antioxidant assays demonstrated significant reductions in intracellular ROS and lipid peroxidation, key markers of oxidative damage. The ability of fruit extracts to activate the Nrf2/HO-1 pathway, a master regulator of cellular antioxidant defenses, provides a plausible mechanistic explanation for these effects. Enhanced expression of Nrf2 and HO-1 following combined treatments indicates that synergistic interactions potentiate the induction of endogenous protective mechanisms, beyond what single extracts achieve. This is crucial since chronic oxidative stress contributes to the pathogenesis of cardiovascular diseases, neurodegeneration, and cancer [refs].

In immune cells, fruit extracts substantially reduced LPS-induced pro-inflammatory cytokine secretion, including TNF- α , IL-6, and IL-1 β . The suppression of NF- κ B p65 nuclear translocation further confirms that these bioactives modulate key inflammatory signaling pathways. The anti-inflammatory potential was strongest in combined treatments, supporting the rationale for dietary intake of diverse fruits to maximize health benefits. Notably, isolated compounds such as quercetin and chlorogenic acid recapitulated these effects, yet combinations often exhibited superior efficacy, highlighting the importance of phytochemical synergy.

The synergy analysis using combination index values consistently showed synergistic interactions for antioxidant and anti-inflammatory endpoints. This underscores the complexity of phytochemical interactions within whole food matrices and the potential advantages of combined therapies or dietary approaches over isolated supplements. Synergism may arise from complementary mechanisms, such as free radical scavenging, metal chelation, enzyme modulation, and signaling pathway interference.

Several limitations of this study should be acknowledged. The *in vitro* cell models, while informative, do not fully replicate the complexity of human physiology, including bioavailability, metabolism, and tissue-specific responses. Future studies involving animal models and clinical trials are necessary to validate the translational relevance of these findings. Moreover, the interactions among bioactives in

complex food matrices warrant further exploration using metabolomics and systems biology approaches.

In conclusion, this research demonstrates that fruit extracts from blueberry, pomegranate, and orange, along with their active components, exert potent protective effects against oxidative stress and inflammation at the cellular level. The synergistic benefits of combined applications provide a strong scientific basis for the promotion of diverse fruit consumption in disease prevention strategies. These findings contribute to the growing body of evidence supporting the use of natural phytochemicals as complementary agents in maintaining cellular health and combating chronic inflammatory conditions.

Conclusion

The current study provides comprehensive insights into the cellular effects of fruit extracts from blueberry, pomegranate, and orange, as well as their principal active components, both individually and in synergistic combinations. Through rigorous *in vitro* experiments using human epithelial and immune cell models, we demonstrated that these fruit-derived bioactives modulate critical cellular processes implicated in oxidative stress and inflammation—two fundamental contributors to a broad spectrum of chronic diseases.

Our findings indicate that moderate concentrations of these extracts enhance cell viability and proliferation, while higher doses exhibit mild cytotoxicity, emphasizing the importance of dosage in therapeutic applications. Notably, combined treatments consistently showed improved safety profiles and more pronounced biological effects, highlighting the value of phytochemical synergy that mimics natural dietary patterns rather than isolated compound supplementation.

Mechanistically, the antioxidant capacity of these extracts is underscored by their activation of the Nrf2/HO-1 signaling pathway, resulting in decreased reactive oxygen species and lipid peroxidation levels. Concurrently, the attenuation of LPS-induced inflammatory cytokines and inhibition of NF- κ B activation in macrophage-like cells illustrate the anti-inflammatory potential of these natural products. The synergy analyses confirmed that the combined application of extracts and their active components yields enhanced protective effects, supporting the concept that complex phytochemical interactions can optimize cellular health outcomes.

This study's implications extend beyond basic cellular biology, advocating for the incorporation of diverse fruit-based bioactives in dietary regimens as a feasible, natural approach to reduce oxidative damage and chronic inflammation. Such nutritional strategies could contribute meaningfully to the prevention and management of cardiovascular, neurodegenerative, metabolic, and inflammatory diseases, aligning with the broader goals of functional nutrition and integrative medicine.

While our *in vitro* findings are promising, future research should address bioavailability, metabolism, and *in vivo* efficacy through animal models and human clinical trials. Further exploration of the molecular interplay among multiple phytochemicals within whole foods will deepen understanding of their therapeutic potential. Overall, this work reinforces the critical role of natural fruit extracts as multi-target agents in promoting cellular resilience and

systemic health, warranting their continued study and application in nutritional science and public health initiatives.

References

1. Agrawal R, Jha S. Antioxidant and anti-inflammatory effects of fruit polyphenols: A review. *J Nutr Biochem*,2019;65:101-113.
<https://doi.org/10.1016/j.jnutbio.2018.10.002>
2. Benzie IF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: The FRAP assay. *Anal Biochem*,1996;239(1):70-76.
<https://doi.org/10.1006/abio.1996.0292>
3. Chen L, Teng H, Xie Z, Huang Q. Mechanisms of fruit polyphenols in regulating inflammation and oxidative stress. *Food Funct*,2020;11(8):6613-6623.
<https://doi.org/10.1039/D0FO01547E>
4. Cho HJ, Kim SH, Kim YS. Synergistic effects of fruit extracts on antioxidant capacity and inflammatory response in human cell lines. *Food Sci Nutr*,2021;9(2):1038-1049.
<https://doi.org/10.1002/fsn3.2086>
5. García F, López JA, Zafra-Stone S. Bioactive compounds in fruits and their effects on health: A review. *Crit Rev Food Sci Nutr*,2018;58(18):3135-3150. <https://doi.org/10.1080/10408398.2017.1378873>
6. Huang D, Ou B, Prior RL. The chemistry behind antioxidant capacity assays. *J Agric Food Chem*,2005;53(6):1841-1856.
<https://doi.org/10.1021/jf030723c>
7. Khan H, Siddiqui S. The impact of flavonoids on oxidative stress and inflammation. *Phytother Res*,2020;34(1):1-10. <https://doi.org/10.1002/ptr.6491>
8. Lee SY, Lee HJ, Park JH. The role of Nrf2 in the antioxidant effects of fruit polyphenols. *Antioxidants (Basel)*,2022;11(3):510.
<https://doi.org/10.3390/antiox11030510>
9. López-García G, Muñoz J. Anti-inflammatory effects of fruit extracts on macrophage cells. *J Med Food*,2017;20(9):893-900.
<https://doi.org/10.1089/jmf.2016.3843>
10. Opara EI, Chohan M. Advances in fruit processing technologies: Antioxidants and anti-inflammatory activities. *Food Technol Biotechnol*,2014;52(3):353-364. <https://doi.org/10.17113/ftb.52.03.14.3530>
11. Wang S, Wang L, Chen X. Synergistic effects of fruit flavonoids on cellular oxidative stress. *J Funct Foods*,2019;58:140-149.
<https://doi.org/10.1016/j.jff.2019.05.022>
12. Xu H, Wang X, Chen Y. Modulation of inflammatory pathways by fruit polyphenols in human immune cells. *Nutrients*,2021;13(5):1501.
<https://doi.org/10.3390/nu13051501>
13. Zhang Q, Zhao X, Zhang J. Protective role of fruit extracts in oxidative stress-related diseases. *Oxid Med Cell Longev*,2017;2017:1-15.
<https://doi.org/10.1155/2017/9346451>